

# Self-dual Smooth Approximations of Convex Functions via the Proximal Average

Heinz H. Bauschke\*, Sarah M. Moffat† and Xianfu Wang‡

March 30, 2010

## Abstract

The proximal average of two convex functions has proven to be a useful tool in convex analysis. In this note, we express Goebel's self-dual smoothing operator in terms of the proximal average, which allows us to give a simple proof of self duality. We also provide a novel self-dual smoothing operator. Both operators are illustrated by smoothing the norm.

**2010 Mathematics Subject Classification:** Primary 26B25; Secondary 26B05, 65D10, 90C25.

**Keywords:** approximation, convex function, Fenchel conjugate, Goebel's smoothing operator, Moreau envelope, proximal average.

## 1 Introduction

Let  $X$  be the standard Euclidean space  $\mathbb{R}^n$ , with inner product  $\langle \cdot, \cdot \rangle$  and induced norm  $\|\cdot\|$ . It will be convenient to set

$$q = \frac{1}{2} \|\cdot\|^2. \quad (1)$$

Now let  $f: X \rightarrow ]-\infty, +\infty]$  be convex, lower semicontinuous, and proper. Since many convex functions are nonsmooth, it is natural to ask: How can one approximate  $f$  with a smooth function?

---

\*Mathematics, Irving K. Barber School, UBC Okanagan, Kelowna, British Columbia V1V 1V7, Canada. E-mail: heinz.bauschke@ubc.ca.

†Mathematics, Irving K. Barber School, UBC Okanagan, Kelowna, British Columbia V1V 1V7, Canada. E-mail: smoffat99@gmail.com.

‡Mathematics, Irving K. Barber School, UBC Okanagan, Kelowna, British Columbia V1V 1V7, Canada. E-mail: shawn.wang@ubc.ca.

The most famous and very useful answer to this question is provided by the *Moreau envelope* [13, 15], which, for  $\lambda > 0$ , is defined by<sup>1</sup>

$$e_\lambda f = f \square \lambda^{-1} \mathbf{q}. \quad (2)$$

It is well known that  $e_\lambda f$  is smooth and that  $\lim_{\lambda \rightarrow 0^+} e_\lambda f = f$  point-wise; see, e.g., [15, Theorem 1.25 and Theorem 2.26].

Let us consider the norm, which is nonsmooth at the origin.

**Example 1.1 (Moreau envelope of the norm)** Let  $\lambda \in ]0, 1[$ , set  $f = \|\cdot\|$ , and denote the closed unit ball by  $C$ . Then, for  $x$  and  $x^*$  in  $X$ , we have<sup>2</sup>

$$e_\lambda f(x) = \begin{cases} \frac{\|x\|^2}{2\lambda}, & \text{if } \|x\| \leq \lambda; \\ \|x\| - \frac{\lambda}{2}, & \text{if } \|x\| > \lambda, \end{cases} \quad (3)$$

$(e_\lambda f)^* = \iota_C + \lambda \mathbf{q}$ , and  $e_\lambda(f^*)(x^*) = (2\lambda)^{-1} \cdot (\max\{0, \|x^*\| - 1\})^2$ . Consequently,  $(e_\lambda f)^* \neq e_\lambda(f^*)$ .

*Proof.* Either a straight-forward computation or [15, Example 11.26(a)] yields

$$f^* = \iota_C. \quad (4)$$

Next, if  $y \in X$ , then

$$e_{1/\lambda} \iota_C(y) = \inf_{c \in C} \lambda \mathbf{q}(y - c) \quad (5)$$

$$= \frac{\lambda}{2} d_C^2(y) \quad (6)$$

$$= \frac{\lambda}{2} \cdot \begin{cases} (\|y\| - 1)^2, & \text{if } \|y\| > 1; \\ 0, & \text{if } \|y\| \leq 1, \end{cases} \quad (7)$$

and thus

$$e_{1/\lambda} \iota_C(x/\lambda) = \frac{\lambda}{2} \cdot \begin{cases} (\|x/\lambda\| - 1)^2, & \text{if } \|x\| > \lambda; \\ 0, & \text{if } \|x\| \leq \lambda. \end{cases} \quad (8)$$

---

<sup>1</sup>The symbol “ $\square$ ” denotes *infimal convolution*:  $(f_1 \square f_2)(x) = \inf_y (f_1(y) + f_2(x - y))$ .

<sup>2</sup> Here  $\iota_C(x) = 0$ , if  $x \in C$ ;  $\iota_C(x) = +\infty$ , if  $x \notin C$  is the *indicator function*,  $f^*(x^*) = \sup_{x \in X} (\langle x, x^* \rangle - f(x))$  is the *Fenchel conjugate* of  $f$ , and  $d_C = \|\cdot\| \square \iota_C$  is the *distance function*.

By [15, Example 11.26(b) on page 495], we obtain

$$e_\lambda f(x) = \frac{1}{\lambda} \mathfrak{q}(x) - e_{1/\lambda} f^*(x/\lambda) \quad (9)$$

$$= \frac{1}{2\lambda} \|x\|^2 - \frac{\lambda}{2} \cdot \begin{cases} \frac{\|x\|^2}{\lambda^2} - \frac{2\|x\|}{\lambda} + 1, & \text{if } \|x\| > \lambda; \\ 0, & \text{if } \|x\| \leq \lambda \end{cases} \quad (10)$$

$$= \begin{cases} \|x\| - \frac{\lambda}{2}, & \text{if } \|x\| > \lambda; \\ \frac{\|x\|^2}{2\lambda}, & \text{if } \|x\| \leq \lambda \end{cases} \quad (11)$$

and  $(e_\lambda f)^* = f^* + \lambda \mathfrak{q} = \iota_C + \lambda \mathfrak{q}$ . Alternatively, one may use [6, Example 2.16], which provides the proximal mapping of  $f$ , and then use the proximal mapping calculus to obtain these results. ■

While the Moreau envelope has many desirable properties, we see from Example 1.1 that the smooth approximation  $e_\lambda f$  is not *self-dual* in the sense that

$$(e_\lambda f)^* \neq e_\lambda(f^*). \quad (12)$$

It is perhaps surprising that self-dual smoothing operators even exist. The first example appears in [9]. Specifically, Goebel defined

$$G_\lambda f = (1 - \lambda^2)e_\lambda f + \lambda \mathfrak{q} \quad (13)$$

and proved that

$$(G_\lambda f)^* = G_\lambda(f^*), \quad (14)$$

i.e., *Fenchel conjugation and Goebel smoothing commute!* For applications of his smoothing operator, see [9].

*The purpose of this note is two-fold. First, we present a different representation of the Goebel smoothing operator which allows us to prove self-duality using the Fenchel conjugation formula for the proximal average. Secondly, the proximal average is also utilized to obtain a novel smoothing operator. Both smoothing operators are computed explicitly for the norm. The formulas derived show that the new smoothing operator is distinct from the one provided by Goebel.*

For  $f_1$  and  $f_2$ , two functions from  $X$  to  $]-\infty, +\infty]$  that are convex, lower semicontinuous and proper, and for two strictly positive convex coefficients ( $\lambda_1 + \lambda_2 = 1$ ), the *proximal average* is defined by

$$\text{pav}(f_1, f_2; \lambda_1, \lambda_2) = (\lambda_1(f_1 + \mathfrak{q})^* + \lambda_2(f_2 + \mathfrak{q})^*)^* - \mathfrak{q}. \quad (15)$$

See [1, 2, 3, 4, 5, 9, 10] for further information and applications of the proximal average. The key property is the *Fenchel conjugation formula*

$$\text{pav}(f_1, f_2; \lambda_1, \lambda_2)^* = \text{pav}(f_1^*, f_2^*; \lambda_1, \lambda_2); \quad (16)$$

see [4, Theorem 6.1], [2, Theorem 4.3], or [1, Theorem 5.1].

We use standard convex analysis calculus and notation as, e.g., in [14, 15, 18]. In Section 2, we consider Goebel's smoothing operator from the proximal-average view point. The new smoothing operator is presented in Section 3.

## 2 The Goebel smoothing operator

**Definition 2.1 (Goebel smoothing operator)** *Let  $f: X \rightarrow ]-\infty, +\infty]$  be convex, lower semicontinuous and proper, and let  $\lambda \in ]0, 1[$ . Then the Goebel smoothing operator [9] is defined by*

$$G_\lambda f = (1 - \lambda^2)e_\lambda f + \lambda q. \quad (17)$$

Note that (17) and standard properties of the Moreau envelope imply that point-wise

$$\lim_{\lambda \rightarrow 0^+} G_\lambda f = f \quad (18)$$

and that each  $G_\lambda f$  is smooth.

Our first main result provides two alternative descriptions of the Goebel smoothing operator. The first description, item (i) in Theorem 2.2, shows a pleasing reformulation in terms of the proximal average. The second description, item (ii) in Theorem 2.2 is less appealing but has the advantage of providing a simple proof of the *self-duality* (iii) observed by Goebel.

**Theorem 2.2** *Let  $f: X \rightarrow ]-\infty, +\infty]$  be convex, lower semicontinuous and proper, and let  $\lambda \in ]0, 1[$ . Then the following hold<sup>3</sup>.*

- (i)  $G_\lambda f = (1 + \lambda) \text{pav}(f, 0; 1 - \lambda, \lambda) + \lambda q.$
- (ii)  $G_\lambda f = (1 + \lambda)^2 \text{pav}\left(f, q; \frac{1-\lambda}{1+\lambda}, \frac{2\lambda}{1+\lambda}\right) \circ (1 + \lambda)^{-1} \text{Id}.$
- (iii) **(Goebel)**  $(G_\lambda f)^* = G_\lambda(f^*).$

---

<sup>3</sup>Here  $\text{Id}: X \rightarrow X: x \mapsto x$  is the *identity operator*.

*Proof.* Let  $x \in X$ . Then, using (15) and standard convex calculus, we obtain

$$\left( (1+\lambda)^2 \text{pav} \left( f, \mathbf{q}; \frac{1-\lambda}{1+\lambda}, \frac{2\lambda}{1+\lambda} \right) \circ (1+\lambda)^{-1} \text{Id} \right) (x) \quad (19)$$

$$= (1+\lambda)^2 \left( \left( \frac{1-\lambda}{1+\lambda} (f + \mathbf{q})^* + \frac{2\lambda}{1+\lambda} (\mathbf{q} + \mathbf{q})^* \right)^* - \mathbf{q} \right) \left( \frac{x}{1+\lambda} \right) \quad (20)$$

$$= (1+\lambda)^2 \left( \frac{1-\lambda}{1+\lambda} (f + \mathbf{q})^* + \frac{\lambda}{1+\lambda} \mathbf{q} \right)^* \left( \frac{x}{1+\lambda} \right) - \mathbf{q}(x) \quad (21)$$

$$= (1+\lambda) \left( (1-\lambda)(f + \mathbf{q})^* + \lambda \mathbf{q} \right)^* (x) - \mathbf{q}(x) \quad (22)$$

$$= (1+\lambda) \left( \left( (1-\lambda)(f + \mathbf{q})^* + \lambda(0 + \mathbf{q})^* \right)^* - \mathbf{q} \right) (x) + \lambda \mathbf{q}(x) \quad (23)$$

$$= \left( (1+\lambda) \text{pav} (f, 0; 1-\lambda, \lambda) + \lambda \mathbf{q} \right) (x). \quad (24)$$

We have verified that (22) as well as the right sides of (i) and (ii) coincide. Starting from (22) and again applying standard convex calculus, we see that

$$(1+\lambda) \left( (1-\lambda)(f + \mathbf{q})^* + \lambda \mathbf{q} \right)^* (x) - \mathbf{q}(x) \quad (25)$$

$$= (1+\lambda) \left( \left( (1-\lambda)(f + \mathbf{q})^* \right)^* \square (\lambda \mathbf{q})^* \right) (x) - \mathbf{q}(x) \quad (26)$$

$$= (1+\lambda) \left( (1-\lambda)(f + \mathbf{q}) \left( \frac{\cdot}{1-\lambda} \right) \square \frac{1}{\lambda} \mathbf{q} \right) (x) - \mathbf{q}(x) \quad (27)$$

$$= (1+\lambda) \inf_y \left( (1-\lambda)(f + \mathbf{q}) \left( \frac{y}{1-\lambda} \right) + \frac{1}{\lambda} \mathbf{q}(x-y) \right) - \mathbf{q}(x) \quad (28)$$

$$= (1+\lambda) \inf_y \left( (1-\lambda)f \left( \frac{y}{1-\lambda} \right) + (1-\lambda)\mathbf{q} \left( \frac{y}{1-\lambda} \right) + \frac{1}{\lambda} \mathbf{q}(x-y) - \frac{1}{1+\lambda} \mathbf{q}(x) \right) \quad (29)$$

$$= (1-\lambda^2) \inf_y \left( f \left( \frac{y}{1-\lambda} \right) + \mathbf{q} \left( \frac{y}{1-\lambda} \right) + \frac{1}{\lambda(1-\lambda)} \mathbf{q}(x-y) - \frac{1}{1-\lambda^2} \mathbf{q}(x) \right). \quad (30)$$

Simple algebra shows that for every  $y \in X$ ,

$$\mathbf{q} \left( \frac{y}{1-\lambda} \right) + \frac{1}{\lambda(1-\lambda)} \mathbf{q}(x-y) - \frac{1}{1-\lambda^2} \mathbf{q}(x) = \frac{1}{\lambda} \mathbf{q} \left( x - \frac{y}{1-\lambda} \right) + \frac{\lambda}{1-\lambda^2} \mathbf{q}(x). \quad (31)$$

Therefore,

$$(1 + \lambda) \left( (1 - \lambda)(f + \mathfrak{q})^* + \lambda \mathfrak{q} \right)^* (x) - \mathfrak{q}(x) \quad (32)$$

$$= (1 - \lambda^2) \inf_y \left( f\left(\frac{y}{1 - \lambda}\right) + \mathfrak{q}\left(\frac{y}{1 - \lambda}\right) + \frac{1}{\lambda(1 - \lambda)} \mathfrak{q}(x - y) - \frac{1}{1 - \lambda^2} \mathfrak{q}(x) \right) \quad (33)$$

$$= (1 - \lambda^2) \inf_y \left( f\left(\frac{y}{1 - \lambda}\right) + \frac{1}{\lambda} \mathfrak{q}\left(x - \frac{y}{1 - \lambda}\right) + \frac{\lambda}{1 - \lambda^2} \mathfrak{q}(x) \right) \quad (34)$$

$$= (1 - \lambda^2) \inf_z \left( f(z) + \frac{1}{\lambda} \mathfrak{q}(x - z) + \frac{\lambda}{1 - \lambda^2} \mathfrak{q}(x) \right) \quad (35)$$

$$= ((1 - \lambda^2) e_\lambda f + \lambda \mathfrak{q})(x) \quad (36)$$

$$= G_\lambda f(x), \quad (37)$$

which completes the proof of (i) and (ii).

(iii): In view of the conjugate formula  $(\beta^2 h \circ (\beta^{-1} \text{Id}))^* = \beta^2 h^* \circ (\beta^{-1} \text{Id})$ , (ii), and (16), we obtain

$$(G_\lambda f)^* = \left( (1 + \lambda)^2 \text{pav} \left( f, \mathfrak{q}; \frac{1 - \lambda}{1 + \lambda}, \frac{2\lambda}{1 + \lambda} \right) \circ (1 + \lambda)^{-1} \text{Id} \right)^* \quad (38)$$

$$= (1 + \lambda)^2 \left( \text{pav} \left( f, \mathfrak{q}; \frac{1 - \lambda}{1 + \lambda}, \frac{2\lambda}{1 + \lambda} \right) \right)^* \circ (1 + \lambda)^{-1} \text{Id} \quad (39)$$

$$= (1 + \lambda)^2 \text{pav} \left( f^*, \mathfrak{q}^*; \frac{1 - \lambda}{1 + \lambda}, \frac{2\lambda}{1 + \lambda} \right) \circ (1 + \lambda)^{-1} \text{Id} \quad (40)$$

$$= (1 + \lambda)^2 \text{pav} \left( f^*, \mathfrak{q}; \frac{1 - \lambda}{1 + \lambda}, \frac{2\lambda}{1 + \lambda} \right) \circ (1 + \lambda)^{-1} \text{Id} \quad (41)$$

$$= G_\lambda (f^*). \quad (42)$$

The proof is complete. ■

**Remark 2.3** Theorem 2.2(i)&(ii) gives two representations of the Goebel smoothing operator in terms of the proximal average. Goebel [8] discovered a converse formula, which we state next without proof:

$$\text{pav}(f, \mathfrak{q}; \lambda, 1 - \lambda) = \frac{(2 - \lambda)^2}{4} G_{\lambda/(2 - \lambda)} f \circ \left( \frac{2}{2 - \lambda} \text{Id} \right). \quad (43)$$

**Example 2.4** Let  $\lambda \in ]0, 1[$  and set  $f = \|\cdot\|$ . Then, for every  $x \in X$ ,

$$G_\lambda f(x) = \begin{cases} \frac{\|x\|^2}{2\lambda}, & \text{if } \|x\| \leq \lambda; \\ \frac{\lambda\|x\|^2}{2} + (1 - \lambda^2)\|x\| - \frac{\lambda(1 - \lambda^2)}{2}, & \text{if } \|x\| > \lambda. \end{cases} \quad (44)$$

*Proof.* Combine (17) and (3). ■

### 3 A new smoothing operator

We now provide a novel smoothing operator that has a very simple expression in terms of the proximal average.

**Definition 3.1 (new smoothing operator)** Let  $f: X \rightarrow ]-\infty, +\infty]$  be convex, lower semicontinuous and proper, and let  $\lambda \in ]0, 1[$ . Then the  $S_\lambda f$  is defined by

$$S_\lambda f = \text{pav}(f, \mathbf{q}; 1 - \lambda, \lambda). \quad (45)$$

**Theorem 3.2** Let  $f: X \rightarrow ]-\infty, +\infty]$  be convex, lower semicontinuous and proper, and let  $\lambda \in ]0, 1[$ . Set  $\mu = \lambda/(2 - \lambda)$ . Then the following hold.

$$(i) \quad S_\lambda f = (1 - \lambda)e_\mu f \circ \left(\frac{2}{2 - \lambda} \text{Id}\right) + \mu \mathbf{q}.$$

$$(ii) \quad (S_\lambda f)^* = S_\lambda(f^*).$$

*Proof.* (i): Let  $x \in X$ . Then, using (45), (15) and standard convex calculus, we obtain

$$(S_\lambda f)(x) = ((1 - \lambda)(f + \mathbf{q})^* + \lambda(\mathbf{q} + \mathbf{q})^*)^*(x) - \mathbf{q}(x) \quad (46)$$

$$= ((1 - \lambda)(f + \mathbf{q})^* + \frac{\lambda}{2}\mathbf{q})^*(x) - \mathbf{q}(x) \quad (47)$$

$$= \left( (1 - \lambda)(f + \mathbf{q}) \left( \frac{\cdot}{1 - \lambda} \right) \square \frac{2}{\lambda} \mathbf{q} \right)(x) - \mathbf{q}(x) \quad (48)$$

$$= \inf_y \left( (1 - \lambda)f\left(\frac{y}{1 - \lambda}\right) + (1 - \lambda)\mathbf{q}\left(\frac{y}{1 - \lambda}\right) + \frac{2}{\lambda}\mathbf{q}(x - y) - \mathbf{q}(x) \right) \quad (49)$$

$$= (1 - \lambda) \inf_y \left( f\left(\frac{y}{1 - \lambda}\right) + \mathbf{q}\left(\frac{y}{1 - \lambda}\right) + \frac{2}{\lambda(1 - \lambda)}\mathbf{q}(x - y) - \frac{1}{1 - \lambda}\mathbf{q}(x) \right). \quad (50)$$

Simple algebra shows that for every  $y \in X$ ,

$$\mathfrak{q}\left(\frac{y}{1-\lambda}\right) + \frac{2}{\lambda(1-\lambda)}\mathfrak{q}(x-y) - \frac{1}{1-\lambda}\mathfrak{q}(x) = \frac{2-\lambda}{\lambda}q\left(\frac{2x}{2-\lambda} - \frac{y}{1-\lambda}\right) + \frac{\lambda}{(1-\lambda)(2-\lambda)}\mathfrak{q}(x). \quad (51)$$

Therefore,

$$(S_\lambda f)(x) = (1-\lambda) \inf_y \left( f\left(\frac{y}{1-\lambda}\right) + \frac{2-\lambda}{\lambda}q\left(\frac{2x}{2-\lambda} - \frac{y}{1-\lambda}\right) + \frac{\lambda}{(1-\lambda)(2-\lambda)}\mathfrak{q}(x) \right) \quad (52)$$

$$= (1-\lambda) \inf_z \left( f(z) + \frac{2-\lambda}{\lambda}q\left(\frac{2x}{2-\lambda} - z\right) \right) + \frac{\lambda}{2-\lambda}\mathfrak{q}(x) \quad (53)$$

$$= (1-\lambda) \left( f \square_{\frac{1}{\mu}} \mathfrak{q} \right) \left( \frac{2x}{2-\lambda} \right) + \mu \mathfrak{q}(x), \quad (54)$$

as claimed.

(ii): Using (45) and (16), we get

$$(S_\lambda f)^* = (\text{pav}(f, \mathfrak{q}; 1-\lambda, \lambda))^* = \text{pav}(f^*, \mathfrak{q}^*; 1-\lambda, \lambda) = \text{pav}(f^*, \mathfrak{q}; 1-\lambda, \lambda) = S_\lambda(f^*). \quad (55)$$

The proof is complete.  $\blacksquare$

Note that Theorem 3.2(i) and standard properties of the Moreau envelope imply that point-wise

$$\lim_{\lambda \rightarrow 0^+} S_\lambda f = f \quad (56)$$

and that each  $S_\lambda f$  is smooth.

**Example 3.3** Let  $\lambda \in ]0, 1[$  and set  $f = \|\cdot\|$ . Then, for every  $x \in X$ ,

$$S_\lambda f(x) = \begin{cases} \frac{(2-\lambda)\|x\|^2}{2\lambda}, & \text{if } \|x\| \leq \frac{\lambda}{2}; \\ \frac{\lambda\|x\|^2}{2(2-\lambda)} + \frac{2(1-\lambda)}{2-\lambda}\|x\| - \frac{\lambda(1-\lambda)}{2(2-\lambda)}, & \text{if } \|x\| > \frac{\lambda}{2}. \end{cases} \quad (57)$$

*Proof.* Combine (3) and Theorem 3.2(i).  $\blacksquare$

**Remark 3.4** Let  $f = \|\cdot\|$ . The explicit formulas provided in Example 2.4 and Example 3.3 imply that  $G_\alpha f \neq S_\beta f$ , for *all*  $\alpha$  and  $\beta$  in  $]0, 1[$ . Thus, the smoothing operator defined by (45) is indeed new and different from Goebel's smoothing operator.

**Remark 3.5** Given a more complicated function  $f$ , the explicit computation of the smoothing operators  $G_\lambda f$  and  $S_\lambda f$  may not be so easy. However, computational convex analysis provides tools [11, 12] to compute the Moreau envelope numerically which — due to the Moreau envelope formulations (17) and Theorem 3.2(i) — makes it possible to compute the smoothing operators  $G_\lambda f$  and  $S_\lambda f$  numerically.

Finally, other approaches to smooth approximation are: Ghomi's integral convolution method [7], Seeger's ball rolling technique [16], and Teboulle's entropic proximal mappings [17].



## Acknowledgment

Heinz Bauschke was partially supported by the Natural Sciences and Engineering Research Council of Canada and by the Canada Research Chair Program. Sarah Moffat was partially supported by the Natural Sciences and Engineering Research Council of Canada. Xianfu Wang was partially supported by the Natural Sciences and Engineering Research Council of Canada.

## References

- [1] H.H. Bauschke, R. Goebel, Y. Lucet and X. Wang: “The proximal average: basic theory”, *SIAM Journal on Optimization* 19, pp. 766–785, 2008.
- [2] H.H. Bauschke, Y. Lucet and M. Trienis: “How to transform one convex function continuously into another”, *SIAM Review* 50, pp. 115–132, 2008.
- [3] H.H. Bauschke, Y. Lucet and X. Wang: “Primal-dual symmetric antiderivatives for cyclically monotone operators”, *SIAM Journal on Control and Optimization* 46, pp. 2031–2051, 2007.
- [4] H.H. Bauschke, E. Matoušková, and S. Reich: “Projection and proximal point methods: convergence results and counterexamples”, *Nonlinear Analysis* 56, pp. 715–738, 2004.
- [5] H.H. Bauschke and X. Wang: “The kernel average for two convex functions and its applications to the extension and representation of monotone operators”, *Transactions of the American Mathematical Society* 361, pp. 5947–5965, 2009.
- [6] P.L. Combettes and V.R. Wajs: “Signal recovery by proximal forward-backward splitting”, *Multiscale Modeling and Simulation* 4, pp. 1168–1200, 2005.
- [7] M. Ghomi: “The problem of optimal smoothing for convex functions”, *Proceedings of the American Mathematical Society* 130, pp. 2255–2259, 2002.
- [8] R. Goebel, personal communication, 2006.
- [9] R. Goebel: “Self-dual smoothing of convex and saddle functions”, *Journal of Convex Analysis* 15, pp. 179–190, 2008.
- [10] R. Goebel: “The proximal average for saddle functions and its symmetry properties with respect to partial and saddle conjugacy”, *Journal of Nonlinear Convex Analysis*, in press.
- [11] Y. Lucet: “Faster than the fast Legendre transform, the linear-time Legendre transform”, *Numerical Algorithms* 16, pp. 171–185, 1997.
- [12] Y. Lucet, H.H. Bauschke, and M. Trienis: “The piecewise linear-quadratic model for computational convex analysis”, *Computational Optimization and Applications* 43, pp. 95–118, 2009.

- [13] J.J. Moreau: “Proximité et dualité dans un espace hilbertien”, *Bulletin de la Société Mathématique de France* 93, pp. 273—299, 1965.
- [14] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, 1970.
- [15] R.T. Rockafellar and R.J-B Wets, *Variational Analysis*, corrected 3rd printing, Springer-Verlag, 2009.
- [16] A. Seeger: “Smoothing a nondifferentiable convex function: the technique of the rolling ball”, *Revista de Matemáticas Aplicadas* 18, pp. 259–268, 1997.
- [17] M. Teboulle: “Entropic proximal mappings with applications to nonlinear programming”, *Mathematics of Operations Research* 17, pp. 670–690, 1992.
- [18] C. Zălinescu, *Convex Analysis in General Vector Spaces*, *World Scientific Publishing*, 2002.